

## REFERENCES

- Alberdi, M.T., Caloi, L., & Palombo, M.R. (1998). Large mammal associations from the Early Pleistocene: Italy and Spain. In Th. Van Kolfschoten & Gibbard, P.L. (Eds.), *The Dawn of the Quaternary* (pp. 521-532). Proceedings of the SEQS-EuroMam symposium 1996. Medelingen Nederlands Instituut voor Toegepaste Geowetenschappen TNO.
- Alberdi, M.-T., & Ruiz-Bustos, A. (1989). Taxonomia y Biostratigrafia de Equidae (Mammalia, Perissodactyla) en la Cuenca de Guadix-Baza (Granada). *Geologia y Paleontologia de la Cuenca de Guadix-Baza, Trabajos sobre el Neogeno-Cuaternario 11*, 239-270.
- Athanassiou, A., 2001. New data on the *Equus stenonis* from the Late Pliocene locality of Sesklo (Thessaly, Greece). *Géodiversitas 23*, 439-469.
- Aouadi, N. (1999). Etude préliminaire des restes crâniens de Chevaux villafranchiens (Ceysaguet, Haute-Loire). *Bulletin du Musée d'Anthropologie préhistorique de Monaco 40*, 23-42.
- Azzaroli, A. (1990). The genus *Equus* in Europe. *European Neogene Mammal Chronology*, 339-355.
- Azzaroli, A., & Voorhies, M. (1993). The Genus *Equus* in North America. The Blancan species. *Palaeontographia Italica 80*, 175-198.
- Azzaroli, A., Colalongo, M.L., Nakagawa, H., Pasini, G., Rio, D., Ruggieri, G., et al. (1997). The Pliocene-Pleistocene boundary in Italy. In Van Couvering, J.A. (Ed.), *The Pleistocene Boundary and the Beginning of the Quaternary*, (pp. 141-155). World and regional Geology 9, Cambridge University Press.
- Bonifay, M.-F., & Brugal, J.-P. (1996). Biogéographie et biostratigraphie des grandes faunes du Pléistocène inférieur et moyen d'Europe du Sud : apport des gisements français. *Paléo 8*, 19-30.
- Debard, E., Faure, M., & Guérin, C. (1994). Stratigraphie du gisement villafranchien moyen de Saint-Vallier (Drôme). *C.R. Acad. Sci. 318*, 1283-1286.
- De Giuli, C. (1972). On the type form of *Equus stenonis* Cocchi. *Palaeontographia italica 68*, 35-49.
- Eisenmann, V. (1981). Etude des dents jugales inférieures des *Equus* actuels et fossiles. *Palaeovertebrata 10*, 127-226.
- Eisenmann, V. (1985). Les Equidés des gisements de la vallée de l'Omo en Ethiopie (collections françaises). In Coppens, Y., & Clark Howell, F. (Eds.), *Les Faunes plio-pléistocènes de la basse vallée de l'Omo (Ethiopie), T. 1: Périssodactyles, Artiodactyles (Bovidae)* (pp. 13-55). Cahiers de Paléontologie.
- Eisenmann, V. (1986). Comparative osteology of modern and fossil Horses, Halfasses and Asses. In Meadow, & R.H., Uerpman, H.P. (Eds.), *Equids in the ancient world* (pp. 67-116). Beihefte zum Tübinger Atlas des Vorderen Orients, Reihe A, Wiesbaden,.
- Eisenmann, V. & Baylac, M. (2000). Extant and fossil *Equus* (Mammalia, Perissodactyla) skulls: a morphometric definition of the subgenus *Equus*. *Zoologica Scripta 29*, 89-100.
- Eisenmann, V. (2002). The primitive horses of the Vatera Formation (Lesbos, Greece). Proceedings of the 1st International Workshop "On Late Plio/Pleistocene extinction and evolution in the Palearctic. The Vatera site". *Annales Géologiques des Pays Helléniques, 1ère série, volume XXXIX (A)*, 131-153.
- Eisenmann, V. (2004). Les Equidés (Mammalia, Perissodactyla) de Saint-Vallier (Drôme, France) et les Equidés Plio-Pleistocènes d'Europe. In Faure, M., & Guérin, C. (Eds.), *Le gisement pliocène final de Saint-Vallier (Drôme, France)*, (S279-S305). Géobios 37, Mémoire spécial 26.
- Eisenmann, V., & Deng, T. (2005). *Equus qingyangensis* (Equidae, Perissodactyla) of the Lower Pleistocene of Bajiazui, China: evidence for the North American origin of an Old World lineage distinct from *E. stenonis*. Les Ongulés Holarctiques du Pliocène et du Pléistocène, Actes du Colloque international d'Avignon, 19-22 Septembre 2000, *Quaternaire, hors-série 2-2005*, 113-122.
- Eisenmann, V. (2006). Pliocene and Pleistocene Equids: Paleontology versus Molecular Biology. In: Kahlke, R.-D., Maul, L. C. & Mazza, P. (Eds.): Late Neogene and Quaternary biodiversity and evolution:

- Regional developments and interregional correlations. Proceedings volume of the 18th International Senckenberg Conference (VI International Palaeontological Colloquium in Weimar), 25th-20th April 2004. Courier Forschungsinstitut Senckenberg (CFS), 256:71-89, 21 figs, 2pls.
- Eisenmann, V., & Vasiliev, S. (2011). Unexpected finding of a new *Equus* species (Mammalia, Perissodactyla) belonging to a supposedly extinct subgenus in late Pleistocene deposits of Khakassia (southwestern Siberia). *Geodiversitas* 33(3): 519-530.
- Forsten, A., & Eisenmann, V. (1995). *Equus (Plesippus) simplicidens* (Cope), not *Dolichohippus*. *Mammalia* 59, 85-89.
- Forsten, A. (1998). The fossil horses (Equidae, Mammalia) from the Plio-Pleistocene of Liventsovka near Rostov-Don, Russia. *Géobios* 31, 645-657.
- Gabunia, M.K., & Vekua, A.K. (1995). A Plio-Pleistocene hominid from Dmanisi, East Georgia, Caucasus. *Nature* 373, 509-512.
- Heintz, E. (1970). Les Cervidés villafranchiens de France et d'Espagne. *Mémoires du Muséum national d'Histoire naturelle, Nouvelle Série, Série C, Tome XXII*, 2 volumes : 303 et 206 pp.
- Hemmer, H., Kahlke, R.-D., & Vekua, A. K. (2001). The Jaguar - *Panthera onca gombaszoegensis* (Kretzoi, 1938) (Carnivora: Felidae) in the late Lower Pleistocene of Akhalkalaki (South Georgia; Transcaucasia) and its evolutionary and ecological significance. *Geobios* 34-4, 475-486.
- Kahlke, R.-D., 2006, (erschienen 2007). Untermassfeld - A late Early Pleistocene (Epivillafranchian) fossil site near Meiningen (Thuringia, Germany) and its position in the development of the European mammal fauna. - *BAR British Archaeological Reports, International Series 1578*, pp.141.
- Kostopoulos, D.S., & Sen, S. (1999). Late Pliocene (Villafranchian) mammals from Sarikol Tepe, Ankara, Turkey. *Mitteilungen der Bayerischen Staatssammlung für Paläontologie und historische Geologie* 39, 165-202.
- Koufos, G. (1992). Early Pleistocene equids from Mygdonia basin (Macedonia, Greece). *Palaeontographia Italica* 79, 167-199.
- Koufos, G.D., & Kostopoulos, D.S. (1997). Biochronology and succession of the Plio-Pleistocene macromammalian localities of Greece. In Aguilar, J.-P., Legendre, S., Michaux, J. (Eds), Actes du Congrès BiochroM'97 (pp. 619-634). Mémoires et Travaux E.P.H.E., Inst. Montpellier 21.
- Kretzoi, M. (1938). Die raubtiere von Gombaszög nebst einer Übersicht der Gesamtfaua (Ein Beitrag zur Stratigraphie des Altquartärs). *Annales historico-naturales Musei Nationalis Hungarici* 31, 88-157.
- Lacombat, F., (Ed) (2005). Les grands mammifères fossiles du Velay. Les collections paléontologiques du Plio-Pléistocène du Musée Crozatier, Le Puy-en-Velay. *Annales de la Société des Amis du Musée Crozatier, Octobre 2005, 13-14*, 208 pp.
- Maul, L.C., Heinrich, W.-D., Parfitt, S. A., & Paunescu, A.-C. (2007). Comment on the correlation between magnetostratigraphy and the evolution of *Microtus* (Arvicolidae, Rodentia, Mammalia) during the Early and early Middle Pleistocene. In Kahlke, R.-D., Maul, L. C., & Mazza, P. P. A. (Eds.). Late Neogene and Quaternary biodiversity and evolution: Regional developments and interregional correlations. Proceedings of the 18th International Senckenberg Conference (VI International Palaeontological Colloquium in Weimar). Volume II. – *Courier Forschungsinstitut Senckenberg* 259, 243-263. Frankfurt am Main.
- Mausser, M. (1987). Geologische und Palaeontologische Untersuchungen an der altpleistozänen Säugetier-Fundstelle Würzburg-Schalksberg. *Münchener Geowissenschaftliche Abhandlungen, Reihe A, 11*, 77pp.
- Meadow, R.H. (1986). Some Equid Remains from Cayönü, Southeastern Turkey. In Meadow, R.H., & Uerpmann, H.P. (Eds.), *Equids in the ancient world* (pp. 266-301). Beihefte zum Tübinger Atlas des Vorderen Orients, Reihe A, Wiesbaden.
- Meadow, R. (1999). The use of size index scaling techniques for research on archaeozoological collections from the Middle East. *Historia Animalium ex Ossibus. Festschrift für Angela von den Driesch zum 65.*

- Geburtstag. *Internationale Archäologie, Bd 8: Studia honoraria*, 285-300. Verlag Marie Leidorf, Rahden, Westf.
- Nomade, S., Pastre, J.F., Guillou, H., Faure, M., Guérin, C., Delson, E., Debard, E., Voinchet, P., Messenger, E. (2013).  $^{40}\text{Ar}/^{39}\text{Ar}$  constraints on some French landmark Late Pliocene to Early Pleistocene large mammalian paleofaunas: Paleoenvironmental and paleoecological implications. *Quaternary Geochronology* 14,2-15.
- Radulesco, C., & Samson, P. (1990). The Plio-Pleistocene Mammalian succession of the Oltets Valley, Dacic Basin, Romania. *Quartärpaläontologie* 8, 225-232.
- Samson, P. (1975). Les Equidés fossiles de Roumanie. *Geologica Romana* 14, 165-352.
- Sher, A.V. (1987). Olyorian land mammal age of Northeastern Siberia. *Palaeontographia italica* 74, 97-112.
- Sotnikova, M.V. (1989). Khishchnye Mlekopitaiushchie Pliotsena - Rannego Pleistotsena. *Geologicheskij Institut, Trudy* 440, 121pp. Izd. Nauka, Moskva.
- Torre, D. (1987). Pliocene and Pleistocene marine-continental correlations. Proceedings of the VIIIth RCMNS Congress, *Ann. Inst. Geol. Publ. Hung.* 30, 71-77.
- Torre, D., Albanelli, A., Bertini, A., Ficarelli, G., Masini, F., & Napoleone, G. (1996). Paleomagnetic calibration of Plio-Pleistocene mammal localities in central Italy. *Acta zool. cracov.* 3, 559-570.
- Uerpmann, H.P. (1982). Faunal remains from Shams ed-din Tannira, a Halafian Site in Northern Syria. *Beyrutus* 30, 3-52.
- Van den Meulen, A.J., & Van Kolfschoten, Th. (1986). Review of the late Turolian to early Biharian mammal faunas from Greece and Turkey. *Memorie della Societa Geologica Italiana* 31, 201-211.
- Vekua, A.K. (1986). The Lower Pleistocene Mammalian Fauna of Akhalkalaki (Southern Georgia, USSR). *Palaeontographia Italica* 74, 63-96.
- Westerhoff, W.E., Cleveringa, P., Meijer, T., van Kolfschoten, TH., & Zagwijn, W.H. (1998). The Lower Pleistocene fluvial (clay) deposits in the Maalbeek pit near Tegelen. In Th. Van Kolfschoten, & Th., Gibbard, P.L. (Eds.), *The Dawn of the Quaternary* (pp. 35-69). Proceedings of the SEQS-EuroMam symposium 1996. Medelingen Nederlands Instituut voor Toegepaste Geowetenschappen TNO.